Mental Health and Substance Use During COVID-19

Summary Report

Conducted by **Leger** for the

Canadian Centre on Substance Use and Addiction and the Mental Health Commission of Canada







October 13 – November 2, 2020 (Time 1) November 19 – December 2, 2020 (Time 2)

Key Findings

- The mental health and substance use impacts of the pandemic have been greater for people living with, or at risk of, mental health and substance use disorders.
 - Only 2 in 5 respondents report strong (very good/excellent) mental health.
 - 14% of respondents report moderately severe/severe current symptoms of depression, 24% report moderate/severe symptoms of anxiety, and 5–6% have seriously contemplated suicide since March 2020.
 - 1 in 3 respondents who use alcohol report increased use and 1 in 5 report problematic use; 2 out of 5 who use cannabis report increased use AND problematic use.
- The pandemic is amplifying the close relationship between mental health and substance use.
 - Respondents with past and current substance use concerns report more mental health symptoms. Nearly 1 out of 2 respondents with past substance use disorders report moderately severe to severe depression symptoms since March 2020.
 - Respondents with past and current mental health concerns report greater increases in substance use. Almost 1 out of 2
 respondents with current mental health symptoms who consume cannabis report increased consumption.
- Access to services is not keeping up with increasing need.
 - Only 22% of respondents with current mental health symptoms and 24% with current problematic substance use report access to treatment since March; about 1 in 5 of respondents who have received care report finding access difficult.

Context and Objectives

Context

The COVID-19 pandemic has increased stress across the population on many fronts. Mental health impacts are increasingly clear, including symptoms of anxiety, depression and suicidal ideation. Alcohol and cannabis use have also gone up. It is well-known that mental health and substance use are intimately connected. Yet, this intersection has not received a great deal of attention during the COVID-19 pandemic. Accordingly, the Mental Health Commission of Canada (MHCC) and the Canadian Centre on Substance Use and Addiction (CCSA) have collaborated in an on-going polling initiative.

This report, which is the first in a series, presents data collected during two timepoints:

- Time 1 (T1): October 13 to November 2, 2020
- Time 2 (T2): November 19 to December 2, 2020

Objectives

The objectives of this project are to:

- Monitor the ongoing impacts of the COVID-19 pandemic on mental health and substance use
- Inform the development of resources to address the mental health and substance use needs of Canadians
- Inform decisions and policies to facilitate greater availability of, and accessibility to, mental health and substance use services during a public health crisis



Lightbulbs are used throughout to provide context for the findings presented in this report. They are not meant to be used as a direct comparison between surveys given differences in methodology. This includes differences in sample characteristics, date of survey, and measurement of mental health and substance use.

Methodology

Study Population

• Residents of Canada, aged 16 and older.

Data Collection

- Survey data was collected at two time periods: between October 13 and November 2, 2020 (Time 1: T1), and between November 19 and December 2, 2020 (Time 2: T2).
- A total of n=2502 (T1) and n=1507 (T2) online surveys were conducted via Leger's online panel (see final slides for sample details).
- This report includes findings for the following priority populations: people reporting any lifetime mental health diagnosis, people reporting any lifetime substance use diagnosis, people reporting current mental health symptoms, and people reporting current problematic alcohol or cannabis use (see next slide details).

Statistical Analysis

- As a non-random online survey, a margin of error is technically not reported.
- If the data were collected through a random sample, the margin of error for T1 would be Canada (n=2502) ±2.0%, 19 times out of 20, and for T2 would be Canada (n=1507) ±2.5%, 19 times out of 20.
- While the sample has been weighted according to age, gender and region using data from the 2016 census, it is not fully representative of the Canadian population and caution should be exercised when comparing results with other surveys.
- The numbers presented have been rounded to the nearest whole number. However, raw values were used to calculate the sums presented and therefore may not correspond to the manual addition of these numbers.


PRIORITY POPULATION PROFILE

| | T1 (Oct/Nov) | T2 (Nov/Dec) |
|---|-----------------|-----------------|
| n = | 2502 | 1507 |
| Lifetime Mental Health Diagnosis [Lifetime MHD] Has a medical or psychological professional ever diagnosed you with a mental health disorder (e.g., depression, anxiety, PTSD, etc.)? YES | 30% (n=747) | 28% (n=428) |
| Lifetime Substance Use Diagnosis [Lifetime SUD] Has a medical or psychological professional ever diagnosed you with substance use disorder (e.g., alcohol use disorder, cannabis use disorder, etc.)? YES | 6% (n=138) | 5% (n=77) |
| Current Mental Health Symptoms [Current MH] Respondents reporting mild to severe symptoms of depression (PHQ-9 score 5+) and/or anxiety (GAD-7 score 5+), and/or recently contemplating suicide. | 59% (n=1482) | 59% (n=883) |
| Current Problematic Substance Use [Current SU] Respondents reporting symptoms of problematic alcohol use (AUDIT score 8+) and/or cannabis use (CUDIT-R score 8+). | 19% (n=465) | 18% (n=265) |

Detailed Results State of Mental Health

About 40% of respondents report strong mental health

Lower % among those with mental health and substance use concerns




Q7: In general, how would you describe your mental health? [same for T1 and T2]

^{*}Statistically significant difference than rest of sample at 95% confidence interval.

14% of respondents report severe depression symptoms

Higher % among those with substance use concerns, past mental health diagnosis



■ 15-19: Moderately severe

■ 10-14: Moderate

■ 5-9: Mild

■ 0-4: None-minimal



According to the Canadian Community Health Survey (retrieved from ODESI), **2%** of Canadians reported moderately severe/severe symptoms of depression in 2015-16.

Q9: DEPRESSION SCORE (PHQ-9) [same for T1 and T2]


(PHQ-9)

^{*}Statistically significant difference than rest of sample at 95% confidence interval.

24% of respondents report moderate to severe anxiety symptoms

Higher % among those with substance use concerns, past mental health diagnosis

Anxiety Symptoms (CAD-7)



Q8: ANXIETY SCORE (GAD-7) [same for T1 and T2]


^{*}Statistically significant difference than rest of sample at 95% confidence interval.

About 5% of respondents have seriously contemplated suicide since March

Higher % among those with substance use concerns, past mental health diagnosis

Suicidal Ideation

Seriously contemplating suicide since March 2020





According to <u>Statistics Canada</u>, **3%** of Canadians in 2019 reported seriously contemplating suicide in the past 12 months.

Q11: Have you seriously contemplated suicide since the COVID-19 pandemic began? [same for T1 and T2]


^{*}Statistically significant difference than rest of sample at 95% confidence interval.

Detailed Results Alcohol and Cannabis Use

About 30% of respondents who use alcohol report using more during pandemic

Higher % among those with mental health and substance use concerns

Change in Alcohol Use




Q22r14: Drinking alcohol (beverages/drinks) - Have you engaged in the following activities more or less since the onset of the COVID-19 pandemic (since March 2020)? [T1] During the past month, have you engaged in more or less of the following activities? [T2].

^{*}Statistically significant difference than rest of sample at 95% confidence interval.

More than 20% of respondents who use alcohol report problematic use

Higher % among those with mental health concerns and a history of substance use disorders




Q30: AUDIT (Alcohol Use Disorder Identification Test) Score [same for T1 and T2] *Statistically significant difference than rest of sample at 95% confidence interval.

About 40% of respondents who use cannabis report using more during pandemic

Higher % among those with mental health and substance use concerns

Change in Cannabis Use




Q22r15: Consuming cannabis - Have you engaged in the following activities more or less since the onset of the COVID-19 pandemic (since March 2020)? [T1] During the past month, have you engaged in more or less of the following activities? [T2]

^{*}Statistically significant difference than rest of sample at 95% confidence interval


About 40% of respondents who use cannabis report problematic use

Higher % among those with mental health concerns and a history of substance use disorders

Problematic Cannabis Use (CUDIT-R)










According to <u>Health Canada</u>, it is estimated that **1 in 11** who use cannabis will develop an addiction to it. If a person smokes cannabis daily, the risk of addiction is **25% to 50%**.

Detailed Results Stress and Coping

Financial situation, social isolation and health top stressors during pandemic




Q18: Thinking about your day-to-day life during the pandemic, what would you say is your biggest source of COVID-19 related stress right now? [same for T1 and T2]

More than 60% of respondents report that they can deal with COVID-19 stress

Lower % of those with mental health and substance use concerns say the same


Dealing with Stress



Q21: When faced with COVID-19 related stress regarding [Q18], you have the personal ability to deal with the situation. Do you...? [same for T1 and T2] *Statistically significant difference than T1 at 95% confidence interval. #Statistically significant difference than rest of sample at 95% confidence interval.


Around 50% of respondents report more online streaming, watching/reading the news and social media since March 2020

Changes in Activities (Since March 2020)




About 30% of respondents report less exercise and social interaction in late November/early December 2020

Changes in Activities (Past month)




Detailed Results Access to Treatment

15% of all respondents and 22% with current mental health symptoms report accessing mental health treatment since March; 1 in 5 found access difficult



Q44: Since March 2020 (T1)/over the past month (T2), have you accessed formal treatment services to help manage your emotions or mental health? Q45: How easy or difficult has it been to access treatment services for mental health? [same for T1 and T2] *Statistically significant difference than rest of sample at 95% confidence interval.

7% of all respondents and 24% with current problematic substance use report accessing substance use treatment since March; 1 in 5 found access difficult



Q42r1,2,3,4: Since March 2020 (T1)/over the past month (T2), have you accessed formal treatment services to help manage your use of [alcohol], [cannabis], [prescription psychoactive drugs], [other illegal psychoactive drugs]? Q45: How easy or difficult has it been to access formal treatment services for substance use? [same for T1 and T2] *Statistically significant difference than rest of sample at 95% confidence interval.



RESPONDENT PROFILE -(1/2)

| | T1 (Oct/Nov) | T2 (Nov/Dec) |
|--------------------------|-----------------|-----------------|
| n = | 2502 | 1507 |
| Gender | (%) | (%) |
| Male | 51 | 51 |
| Female | 48 | 47 |
| Other | | 1 |
| Age | | |
| 16 to 24 | 13 | 13 |
| 25 to 39 | 25 | 24 |
| 40 to 64 | 42 | 42 |
| 65 years of age or older | 21 | 21 |
| 2SLGBTQ+ | | |
| Yes | 11 | 10 |
| No | 88 | 89 |

| | T1 (Oct/Nov) | T2 (Nov/Dec) |
|---|-----------------|-----------------|
| n = | 2502 | 1507 |
| Region | (%) | (%) |
| British Columbia | 13 | 13 |
| Alberta | 12 | 12 |
| Saskatchewan | 3 | 3 |
| Manitoba | 4 | 4 |
| Ontario | 39 | 39 |
| Quebec | 23 | 23 |
| Atlantic | 7 | 7 |
| Canadian Status | | |
| Canadian citizens by birth / non-immigrants | 79 | 79 |
| Canadian by naturalization | 13 | 13 |
| Permanent resident | 6 | 6 |
| No status | 1 | 1 |

RESPONDENT PROFILE -(2/2)

| | T1 (Oct/Nov) | T2 (Nov/Dec) |
|---|-----------------|-----------------|
| n = | 2502 | 1507 |
| Ethnicity | (%) | (%) |
| White | 77 | 76 |
| East/Southeast Asian | 8 | 8 |
| Indigenous Peoples (First Nations, Inuk, Inuit, Métis) | 5 | 5 |
| South Asian | 5 | 5 |
| Black | 2 | 4 |
| Middle Eastern | 2 | 2 |
| Latino | 1 | 2 |
| Canadian / French Canadian (Non-specified) | 1 | <1 |
| Income | | |
| Less than \$35K | 21 | 19 |
| \$35K to \$75K | 30 | 31 |
| \$75K to \$100K | 16 | 16 |
| \$100K and over | 24 | 23 |

| | T1 (Oct/Nov) | T2 (Nov/Dec) |
|------------------------------------|-----------------|-----------------|
| n = | 2502 | 1507 |
| Education | (%) | (%) |
| Less than a high school diploma | 3 | 3 |
| High school degree or equivalent | 8 | 21 |
| Some college, no degree | 31 | 29 |
| Bachelor's degree | 31 | 31 |
| Master's degree | 8 | 7 |
| Professional degree | 6 | 6 |
| Doctorate | 1 | 2 |
| Employment | | |
| Employed | 46 | 45 |
| Unemployed prior to COVID | 5 | 4 |
| Laid off or unemployed since COVID | 5 | 5 |
| Student | 10 | 10 |
| Retired | 24 | 26 |
| Self-employed | 5 | 6 |
| Unable to work | 4 | 4 |

